arXiv:1505.06435v1 [math.CV] 24 May 2015

Galina Bakhtina, Inna Dvorak, Iryna Denega

Problems on extremal decomposition of the complex plane

Considered here is one quite general problem about description of extremal configurations maximizing the product of inner radii non-overlapping domains.

Let $\mathbb{N}$, $\mathbb{R}$ be a set of natural and real numbers, respectively, $\mathbb{C}$ be a complex plane, $\overline{\mathbb{C}} = \mathbb{C}\bigcup\{\infty\}$ be a Riemann sphere and $\mathbb{R}^+ = (0, \infty)$.

Let $r(B, a)$ be a inner radius of domain $B \subset \overline{\mathbb{C}}$, with respect to a point $a \in B$ (see [1, 2, 3, 7, 11]).

Let $n \in \mathbb{N}$. A set of points $A_n := \left\{a_k \in \mathbb{C} : \, k = \overline{1, n}\right\}$, is called *n-radial* system, if $|a_k| \in \mathbb{R}^+$, $k = \overline{1, n}$, and $0 = \arg a_1 < \arg a_2 < \ldots < \arg a_n < 2\pi$.

Denote

$$\alpha_k := \frac{1}{\pi} \arg \frac{a_{k+1}}{a_k}, \; \alpha_{n+1} := \alpha_1, \; k = \overline{1, n}.$$

Consider next functional

$$J_n(\gamma) = r^{\gamma}\left(B_0, 0\right) \prod_{k=1}^{n} r\left(B_k, a_k\right), \tag{1}$$

where $\gamma \in \mathbb{R}^+$, B_0, B_1, B_2,..., B_n, $n \geq 2$ is non-overlapping domains (e.i. $B_p \cap B_j = \emptyset$ if $p \neq j$) in $\overline{\mathbb{C}}$, $a_0 = 0$, $|a_k| = 1$, $k = \overline{1, n}$, $a_k \in B_k$, $k = \overline{0, n}$ and $\gamma \leq n$.

One of the main concepts in this paper is a notion of quadratic differential. Quadratic differential is a convenient tool of describing extremals on the plane. A great collection of definitions and results about quadratic differentials one can find in monographs by well known analyst J. Jenkins [4].

For example, we consider next quadratic differential

$$Q(w)dw^2 = -\frac{(n^2 - \gamma)w^n + \gamma}{w^2(w^n - 1)^2}. \tag{2}$$

If $n = 3$ and $\gamma = 1$ then (2) has four circular domains B_0, B_1, B_2, B_3 such that $\overline{B_0 \cup B_1 \cup B_2 \cup B_3} = \overline{\mathbb{C}}$. A simple poles of the quadratic differential (2) are $w_1 = -\frac{1}{2}$, $w_2 = \frac{1}{4} + \frac{\sqrt{3}}{4}i$, $w_3 = \frac{1}{4} - \frac{\sqrt{3}}{4}i$. The structure of the trajectories of (2) is shown in Fig. 1.



Figure 1: Structure of the trajectories of the quadratic differential (2) for $n = 3$ and $\gamma = 1$.

Let

$$F_\delta(x) = 2^{x^2+6} \cdot x^{x^2+2-2\delta} \cdot (2-x)^{-\frac{1}{2}(2-x)^2}(2+x)^{-\frac{1}{2}(2+x)^2},$$
$$x \in (0, 2], \quad 0 \leqslant \delta \leqslant 0,7.$$

We proved

Theorem. *Let $n \in \mathbb{N}$, $n \geqslant 5$, $\gamma \in (0; 1,75]$, $0 \leqslant \delta \leqslant 0,7$. Then for any n-radial system of points $A_n = \{a_k\}_{k=1}^{n}$, $|a_k| = 1$, and any system of non-overlapping domains B_k, $a_k \in B_k \subset \overline{\mathbb{C}}$, $k = \overline{0,n}$, $a_0 = 0$, we have inequality*

$$J_n(\gamma) \leqslant \gamma^{-\frac{\delta \cdot n}{2}} \cdot \left(\prod_{k=1}^{n} \alpha_k\right)^{\delta} \cdot \left[F_\delta\left(\frac{2}{n}\sqrt{\gamma}\right)\right]^{\frac{n}{2}},$$

where equality holds if a_k and B_k, $k = \overline{0,n}$, are, respectively, poles and circular domains of the quadratic differential (2).

Method of proof of this theorem is based on using of separating transformation [6, 7] and ideas of works [9, 11, 15, 16].

In an analogous way as in papers [15, 16, 17] we obtain next inequality for functional (1)

$$J_n(\gamma) \leqslant \gamma^{-\frac{\delta n}{2}} \left(\prod_{k=1}^{n} \alpha_k\right)^{\delta} \left[\prod_{k=1}^{n} F_\delta\left(\alpha_k\sqrt{\gamma}\right)\right]^{1/2}, \ \delta \in [0; 0,7]. \tag{3}$$

Consider the functional

$$\tilde{J}_n(\gamma) = \gamma^{\frac{\delta n}{2}} \left(\prod_{k=1}^{n} \alpha_k\right)^{-\delta} J_n^{(\gamma)}.$$

It follows from (3) that

$$\tilde{J}_n(\gamma) \leqslant \left[\prod_{k=1}^{n} F_\delta\left(\alpha_k \sqrt{\gamma}\right)\right]^{\frac{1}{2}}.$$

Further, consider an extremal problem

$$\prod_{k=1}^{n} F_\delta(x_k) \longrightarrow \max, \quad \sum_{k=1}^{n} x_k = 2\sqrt{\gamma}, \quad x_k = \alpha_k\sqrt{\gamma}$$

$$0 < x_k \leqslant 2, \quad 0 \leqslant \delta \leqslant 0,7.$$

Let $\Psi_\delta(x) = \ln\left(F_\delta(x)\right)$. And let $X^{(0)} = \left\{x_k^{(0)}\right\}_{k=1}^{n}$ be an arbitrary extremal point of this problem.

It follows from the paper [9] that: if $0 < x_k^{(0)} < x_j^{(0)} < 2$, $k \neq j$, than

$$\Psi'_\delta(x_k^{(0)}) = \Psi'_\delta(x_j^{(0)}),$$

where $k, j = \overline{1, n}$, $k \neq j$, $0 \leqslant \delta \leqslant 0,7$,

$$\Psi'_\delta(x) = 2x\ln(2x) + (2-x)\ln(2-x) - (2+x)\ln(2+x) + \frac{2}{x} - \frac{2\delta}{x}$$

(see Fig. 2).



Figure 2: Graph of the function $\Psi'_\delta(x)$

We will show that the following assertion is true

$$x_1^{(0)} = x_2^{(0)} = \ldots = x_n^{(0)}.$$

Let

$$\sigma_1 := \sigma_1(\delta,\gamma) = \min_{1\leqslant k\leqslant n} x_k^{(0)}(\delta,\gamma),$$

$$\sigma_0 := \sigma_0(\delta,\gamma) = \max_{1\leqslant k\leqslant n} x_k^{(0)}(\delta,\gamma),$$

$\sigma_1 \leqslant \sigma_k \leqslant \sigma_0$, $k=\overline{1,n}$, $0\leqslant\delta\leqslant 0,7$, $\gamma\in(0;1,75]$.

A function $\Psi''_\delta(x)$ is strictly increasing on $(0,2)$ for each fixed δ. Thus

$$sign\Psi''_\delta(x) \equiv sign(x - x_0(\delta,\gamma)).$$

It is not difficult to see if $\sigma_0 \leqslant x_0(\delta,\gamma)$ then from the conditions of the problem we obtain $x_1^{(0)} = \ldots = x_n^{(0)}$.

Assume $x_0(\delta,\gamma)\leqslant\sigma_0 < 1,62$. Then

$$\sigma_1 \leqslant \frac{1}{n-1}\sum_{k=1}^{n-1} x_k^{(0)} = \frac{2\sqrt{\gamma}-\sigma_0}{n-1} \leqslant \frac{2-\sigma_0}{n-1}.$$

Therefore for $n\geqslant 5$, $\gamma\in(0;1,75]$, $0\leqslant\delta\leqslant 0,7$, inequality holds

$$\sigma_1 \leqslant (2\sqrt{\gamma}-\sigma_0)/(n-1) \leqslant (2,645751 - x_0(\delta,\gamma))/4 \leqslant$$

$$\leqslant (2,645751 - 1,08441)/4 < 0,390335.$$

Thus

$$\Psi'_\delta(\sigma_1) > \Psi'_\delta(0,390335) > \Psi'_{0,7}(0,390335) = 0,027069 >$$
$$> 0,018707 = \Psi'_0(1,62) \geqslant \Psi'_\delta(\sigma_0).$$

Hence $\sigma_0 \notin [x_0(\delta,\gamma);\ 1,62)$.

Let $1,62\leqslant\sigma_0\leqslant 2$. Then for $n\geqslant 5$, $\gamma\in(0;1,75]$, $0\leqslant\delta\leqslant 0,7$, we have

$$\sigma_1 \leqslant (2\sqrt{\gamma}-\sigma_0)/(n-1) \leqslant (2,645751 - 1,62)/4 < 0,256438.$$

That is

$$\Psi'_\delta(\sigma_1) > \Psi'_\delta(0,256438) > \Psi'_{0,7}(0,256438) = 1,130326 >$$

$$> 1 = \Psi'_0(2) \geqslant \Psi'_\delta(\sigma_0).$$

So, $\sigma_0 \notin [x_0(\delta,\gamma);\ 2)$.

Therefore, extremal set $\left\{x_k^{(0)}\right\}_{k=1}^{n}$ is possible only if

$$x_1^{(0)} = x_2^{(0)} = \ldots = x_n^{(0)}.$$

From the foregoing the following relation holds

$$\prod_{k=1}^{n} F_\delta\left(\alpha_k\sqrt{\gamma}\right) \leqslant \left[F_\delta\left(\frac{2}{n}\sqrt{\gamma}\right)\right]^n.$$

A sign of equality is verified directly. Theorem is thus proved.

Bibliography

[1] M.A. Lavrent'ev. *On the theory of conformal mappings.* Tr. Sci. Inst An USSR. **5**, 159–245 (1934). (in Russian)

[2] G.M. Goluzin. *Geometric function theory of complex variable.* Moscow: Science. 628 (1966). (in Russian)

[3] V.K. Hayman. *Multivalent functions.* Moscow:Publishing House of Foreign Literature. 180 (1960).

[4] J.A. Jenkins. *Univalent functions and conformal mapping.* Moscow:Publishing House of Foreign Literature. 256 (1962). (in Russian)

[5] P.M. Tamrazov. *Some extremal problems in the theory of univalent conformal mappings.* Math. collection. **67** (109), no.3, 329–337 (1965). (in Russian)

[6] V.N. Dubinin. *Separating transformation of domains and problems on extremal decomposition.* Notes scientific. sem. Leningr. Dep. of Math. Inst. AN USSR. **168**, 48–66 (1988). (in Russian)

[7] V.N. Dubinin. *Symmetrization method in geometric function theory of complex variables.* Successes Mat. Science. **49**, no.1 (295), 3–76 (1994). (in Russian)

[8] L.I. Kolbina. *Conformal mapping of the unit circle onto non-overlapping domains..* Bulletin of Leningrad University. **5**, 37–43 (1955). (in Russian)

[9] L.V. Kovalev. *To the problem of extremal decomposition with free poles on the circle.* Far Eastern Math. collection. **2**, 96–98 (1996). (in Russian)

[10] G.V. Kuzmina. *Extremal metric method in problems of the maximum of product of powers of conformal radii of non-overlapping domains with free parameters.* Notes scientific. sem. Leningr. Dep. of Math. Inst. AN USSR. **302**, 52–67 (2003). (in Russian)

[11] A.K. Bakhtin, G.P. Bakhtina, Yu.B. Zelinskii. *Topological-algebraic structures and geometric methods in complex analysis.* Proceedings of the Institute of Mathematics of NAS of Ukraine. 308 (2008). (in Russian)

[12] V.N. Dubinin. *Capasities of condensers and symmetrization in geometric function theory of complex variables.* Vladivostok: Dal'nayka. 390 (2009). (in Russian)

[13] A.K. Bakhtin, G.P. Bakhtin, R.V. Podvisotskii. *Inequalities for inner radii of non-overlapping domains.* Reports NAS of Ukraine. no.9, 7 — 11 (2009). (in Russian)

[14] Y.V. Zabolotnii. *Application of separation transformation in one problem on non-overlapping domains*. Reports of NAS of Ukraine. no.9, 11 — 14 (2011). (in Russian)

[15] A.K. Bakhtin, I.V. Denega. *Addendum to a theorem on extremal decomposition of the complex plane*. Bulletin de la société des sciences et des lettres de Łódź, Recherches sur les déformations. V. LXII, no.2, 83—92 (2012).

[16] A.K. Bakhtin, I.V. Denega. *Separating tranformation in problems of maximizing the product of powers of the inner radii of non-overlapping domains*. Proceedings of the Institute of Mathematics of NAS of Ukraine. V.9, no.2, 32—44 (2012). (in Russian)

[17] A.K. Bakhtin, I.V. Denega. *About one Dubinin's problem*. Proceedings of the Institute of Mathematics of NAS of Ukraine. V.10, no.4-5, 401—411 (2013). (in Russian)